

A finite dimensional approximation for pricing moving average options.

March 3, 2020

Premia 22

The following method computes the price of American Options whose payoff depends on the moving average of the underlying asset price. It is based on the paper [1].

1 Introduction

The method computes the price of American Options whose payoff depends on the moving average X defined by

$$X_t = \int_{t-\delta}^t S_u du, \quad \forall t \geq \delta$$

where S represents the underlying asset and δ is a fixed time window. The process (S, X) is not Markovian, and in a continuous time framework it is not possible to find n processes (X^1, \dots, X^n) such that (S, X, X^1, \dots, X^n) are jointly Markovian. In a discrete time framework (Bermudan options), n would be equal to the number of time steps within the average window.

The paper proposes a method for pricing moving average American options based on a finite dimensional approximation of the infinite-dimensional dynamics of the moving average process. The approximation is based on a truncated expansion of the weighting measure used for averaging in a series involving Laguerre polynomials, truncated at n terms, which leads to $(n+1)$ -dimensional Markovian approximation to the initial infinite dimensional problem.

2 Theoretical framework

We consider a more general moving average process of the form

$$M_t = \int_0^\infty S_{t-u} \mu(du),$$

where μ is a finite possibly signed measure on $[0, \infty[$. The paper proposes a finite-dimensional approximation to M , that is n processes $X^{p,1}, \dots, X^{p,n}$ such that $(S, X^{p,1}, \dots, X^{p,n})$ are jointly Markov, and M_t is approximated by $M_t^{n,p}$, which depends deterministically on $(S_t, X_t^{p,1}, \dots, X_t^{p,n})$ (see [1, Proposition 2.2]). More precisely, we have

$$M_t^{n,p} = (H(0) - H_n^p(0))S_t + \sum_{k=0}^{n-1} a_k^p X_t^{p,k},$$

where

$$X_t^{p,k} = \int_0^\infty L_k^p(v) S_{t-v} dv, \quad \forall k = 0, \dots, n-1$$

and $L_k^p(t) = \sqrt{2p} P_k(2pt) e^{-pt}$, $k \geq 0$ are the scaled Laguerre functions, in which $(P_k)_{k \geq 0}$ is the family of Laguerre polynomials (see [1, (9)]) and p is a parameter to be fixed. The functions H and H_n^p are defined by

$$H(x) = \mu([x, \infty[), \quad H_n^p(x) = \sum_{k=0}^{n-1} A_k^p L_k^p(x), \quad \text{where } A_k^p = \langle H, L_k^p \rangle.$$

Coefficients $(a_k^p)_k$ are defined in [1, (11)].

2.1 Uniformly weighted measure

When $\mu(dx) = \frac{1}{\delta} \mathbb{1}_{[0, \delta]} dx$, the Laguerre coefficients $A_k^{\delta,p} = \langle H, L_k^p \rangle$ are defined by

$$A_k^{\delta,p} = (-1)^k \frac{\sqrt{2p}}{p} - \frac{1}{p} c_k^{\delta,p} - \frac{2}{p} \sum_{i=0}^{k-1} (-1)^{k-i} c_i^{\delta,p},$$

where

$$c_n^{\delta,p} = \frac{\sqrt{2p}}{\delta p} \left[1 - e^{-p\delta} P_n(2p\delta) + 2 \sum_{k=1}^n (-1)^k (1 - e^{-p\delta} P_{n-k}(2p\delta)) \right].$$

The optimal scale parameter $p_{opt}(\delta, n)$ can be computed using the relation $p_{opt}(\delta, n) = \frac{p_{opt}(1, n)}{\delta}$ and [1, Table 1].

3 Monte Carlo-based numerical method

We consider a uniformly weighted moving average. The following method computes the price of the discrete time version of the American option $\sup_\tau \mathbb{E}[\phi(S_\tau, M_\tau)]$, in which the moving average X has been replaced by its approximation $M^{n, p_{opt}}$, and the exercise is possible on an equidistant time grid π with N time steps $\Delta t = \frac{T}{N}$. The approach corresponds to the one of Longstaff and Schwartz, and the computation of conditional expectations is done with a regression based approach. N_δ denotes the number of time steps within the average window of length δ : $N_\delta = \frac{\delta}{T} N$. The spot price is discretised on π and is written S^π , the discretised version of X is given by

$$X_{t_i}^\pi = \frac{1}{N_\delta} \sum_{j=i-N_\delta+1}^i S_{t_j}^\pi, \quad \forall t_i \in \pi.$$

The discrete time version of the Laguerre processes are defined by

$$X_{t_i}^{p,k,\pi} = \sum_{j=1}^i (S_{t_j}^\pi - S_{t_{j-1}}^\pi) (i-j+1) \Delta t c_k^{(i-j+1)\Delta t, p} + S_0 (-1)^k \frac{\sqrt{2p}}{p}, \quad \forall t_i \in \pi.$$

3.1 The $Lag - LS^*$ algorithm

The backward algorithm works as follows

1. Initialization : $\tau_N^{\pi, (m)} = T, m = 1, \dots, M$
2. Backward induction for $i = N - 1, \dots, N_\delta, m = 1, \dots, M$

$$\begin{aligned}\tau_i^{\pi, (m)} &= t_i \mathbb{1}_{\{A_i^{(m)}\}} + \tau_{i+1}^{\pi, (m)} \mathbb{1}_{\{(A_i^{(m)})^c\}}, \\ A_i^{(m)} &= \left\{ \Phi(S_{t_i}^{\pi, (m)}, X_{t_i}^{\pi, (m)}) \geq \mathbb{E}_{t_i}[\Phi(S_{\tau_{i+1}^\pi}^\pi, X_{\tau_{i+1}^\pi}^\pi)] \right\}\end{aligned}$$

where $\mathbb{E}_{t_i}[\cdot] = \mathbb{E}[\cdot | (S_{t_i}^\pi, X_{t_i}^{popt, 0, \pi}, \dots, X_{t_i}^{popt, n-1, \pi})]$.

Estimators of the conditional expectations are constructed with a Monte Carlo based technique.

3.2 The $NM - LS$ algorithm

The following algorithm is a non-markovian approximation of the previous algorithm. Let $(\theta_i^\pi)_{i=N_\delta, \dots, N}$ denote the discrete time sequence of the estimated optimal exercises times. The algorithm works as follows

1. Initialization : $\theta_N^\pi = T$
2. Backward induction for $i = N - 1, \dots, N_\delta,$

$$\begin{aligned}\theta_i^\pi &= t_i \mathbb{1}_{\{A_i\}} + \theta_{i+1}^\pi \mathbb{1}_{\{(A_i)^c\}}, \\ A_i &= \left\{ \Phi(S_{t_i}^\pi, X_{t_i}^\pi) \geq \mathbb{E}[\Phi(S_{\theta_{i+1}^\pi}^\pi, X_{\theta_{i+1}^\pi}^\pi) | (S_{t_i}^\pi, X_{t_i}^\pi)] \right\}\end{aligned}$$

3. Estimation of the option price at time 0 $U_0^\pi = \mathbb{E}[\Phi(S_{\theta_{N_\delta}^\pi}^\pi, X_{\theta_{N_\delta}^\pi}^\pi)]$.

4 Numerical experiments

4.1 Moving average without time delay

In the case of an American moving average Call, the numerical data used by default are the following

S_0	T	r	σ	r	δ
100	0.2	0.05	0.3	0.05	0.02

The number of trajectories of S is $M = 10^5$. The number of discretization time steps used for the discretization of S is $N = 50$ (then $N_\delta = 5$). The number of polynomial basis functions is 4. When using the $Lag - LS^*$ algorithm, $n = 3$.

4.2 Moving average with time delay

We consider a moving average American option with time delay $l \geq 0$ whose value at time 0 is

$$\sup_{\tau \in \mathcal{T}_{[\delta+l, T]}} \mathbb{E}[\Phi(S_\tau, X_\tau)], \quad X_\tau = \frac{1}{\delta} \int_{\tau-l-\delta}^{\tau-l} S_u du.$$

In this case,

$$X_{t_i}^\pi = \frac{1}{N_\delta} \sum_{j=i-N_\delta-N_l+1}^{i-N_l} S_{t_j}^\pi, \forall t_i \in \pi,$$

where $N_l = l \frac{N}{T}$. The numerical data used by default are the following

S_0	T	r	σ	r	δ	l
100	0.2	0.05	0.3	0.05	0.02	0.1

The number of trajectories of S is $M = 10^5$. The number of discretization time steps used for the discretization of S is $N = 50$ (then $N_\delta = 5$ and $N_l = 25$). The number of polynomial basis functions is 4. When using the *Lag* – *LS** algorithm, $n = 3$.

References

- [1] M. Bernhart, P. Tankov, and X. Warin. A finite dimensional approximation for pricing moving average options. *SIAM Journal of Financial Mathematics*, 2:989–1013, 2011.
[1](#), [2](#)